# Canadian Eating Disorders Strategy

2019-2029

This Strategy has been developed by representatives of the Eating Disorders Association of Canada-Association des Troubles Alimentaires du Canada (EDAC-ATAC), Eating Disorders Foundation of Canada (EDFC), National Eating Disorder Information Centre (NEDIC), and National Initiative for Eating Disorders (NIED). Its purpose is to scan and describe the field of Eating Disorders in Canada and facilitate ongoing dialogue and broad engagement with a range of stakeholders interested in improving health and social outcomes of people living with and affected by EDs in this country.

© The Canadian Eating Disorders Strategy: 2019 - 2029 is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License.

Citation: "Canadian Eating Disorders Alliance (2019). The Canadian Eating Disorders Strategy: 2019 - 2029.

November 2019









# **Table of Contents** Canadian Eating Disorders Strategy: 2019-2029

Preface
Introduction
About the Canadian Eating Disorder
Vision
The Canadian Eating Disorders Strate
Pillar 1: Prevention Pillar 2: Public Education and Av Pillar 3: Treatment Pillar 4: Caregiver Support Pillar 5: Training Pillar 5: Research
Conclusion
How this Strategy Was Develop
Annex 1
Why create a Canadian Eating D
Annex 2
Summary of recommendations
Endnotes

	2
	2
rs Alliance (CEDA)	3
	5
2gy	5
	6
wareness	8
	18
	22
	25
	28
ped	28
	30
Disorders Strategy?	
	34
5	
	42

# Preface

In 2016, EDAC-ATAC, EDFC, NEDIC and NIED<sup>1</sup> joined together to articulate a national strategy for Eating Disorders (EDs). Our four organizations agreed to first describe the field of Eating Disorders in Canada and start a process of creating actions involving many stakeholders that could improve outcomes for people affected by EDs. These actions appear as 'Recommendations' (grouped under 6 pillars) throughout this Strategy.

We welcome the opportunity to discuss these recommendations with stakeholders across Canada to determine how we can best improve health and social outcomes for people living with or recovering from EDs in Canada.

# Introduction

Approximately 1 million<sup>2</sup> Canadians meet the diagnostic criteria for an Eating Disorder (ED), such as Anorexia Nervosa (AN), Bulimia Nervosa (BN), Binge Eating Disorder (BED), Avoidant Restrictive Food Intake Disorder (ARFID)<sup>i</sup> and Otherwise Specified Feeding and Eating Disorder (OSFED)<sup>3</sup>. EDs are serious but treatable mental illnesses that can affect anyone regardless of gender, age, racial and ethnic identity, sexual orientation or socio-economic background<sup>ii</sup>.

Unfortunately, research data indicate that the majority of people with EDs do not seek or receive treatment, and that those who do experience lengthy delays<sup>iii</sup>. Many people with an ED are never diagnosed, do not receive any treatment, and suffer significant personal and family distress. The social and economic costs of untreated EDs are similar to those of depression and anxiety, with debilitating physical and mental health effects comparable to psychosis and schizophrenia<sup>4</sup>.

While it is well documented that mental illnesses are a leading cause of premature death in Canada<sup>iv</sup>, it is less known that EDs have the highest overall mortality rate of any mental illness, with estimates between 10-15%<sup>v</sup>. Suicide is the second leading cause of death (after cardiac disease) among those with AN<sup>vi</sup>; 20% of people with AN and 25-35% of people with BN may attempt suicide in their lifetime<sup>vii</sup>. For females aged 15-24 years old, the mortality rate associated with AN is 12 times greater than that of ALL other causes of death combined<sup>viii</sup>.

Younger Canadians are also increasingly<sup>ix</sup> engaging in dieting behaviour which may put them at risk of developing an ED and other health-compromising conditions<sup>x</sup>. In Canada, between **12% and 30%** of girls and **9% and 25%** of boys aged 10-14 report dieting to lose weight<sup>xi</sup>. Moreover, the incidence of EDs in Canadian children is estimated to be 2 to 4 times greater than Type 2 Diabetes<sup>xii</sup>.

<sup>1</sup> Eating Disorders Association of Canada – Association des Troubles Alimentaires du Canada (EDAC-ATAC); National Eating Disorder Information Centre (NEDIC); Eating Disorders Foundation of Canada (EDFC); and National Initiative for Eating Disorders (NIED).

<sup>2</sup> 2016 calculation based on prevalence rates cited in Langlois, K.A., Samokhvalov, A.V., Rehm, J., Spence, S.T. and Gorber, S.C., *Health Descriptions for Canadians – Mental Illnesses* (2012): 39-42, Statistics Canada surveys and statistical programs – Documentation: 82-619-M2012004, <a href="https://www150.statcan.gc.ca/n1/en/pub/82-619-m/82-619-m/2012004-eng.pdf?st=gRt3lMry">https://www150.statcan.gc.ca/n1/en/pub/82-619-m/82-619-m/2012004-eng.pdf?st=gRt3lMry</a>

<sup>3</sup> In the United States, it is estimated that 30 million Americans live with an ED and that only 1 in 10 receive treatment (National Eating Disorders Association (NEDA), <u>https://www.nationaleatingdisorders.org/statistics-research-eating-disorders</u>). In 2012, it was estimated that there were 913,986 people with an ED in Australia.

<sup>4</sup> The Butterfly Foundation, *Paying the price: The economic and social impacts of eating disorders in Australia* (2012), <u>https://thebutterflyfounda-tion.org.au/about-us/information-and-resources/paying-the-price/</u>



# About the Canadian Eating Disorders Alliance (CEDA)

CEDA is a dedicated group of stakeholders and volunteers whose main goal is to improve health and social outcomes for people living with and affected by EDs. CEDA includes the:

- Eating Disorders Association of Canada-Association des Troubles Alimentaires du Canada (EDAC-ATAC);
- Eating Disorders Foundation of Canada (EDFC);
- National Eating Disorder Information Centre (NEDIC); and
- National Initiative for Eating Disorders (NIED).

#### Purpose

To be the national collective voice for Canadians affected by Eating Disorders, including people living with an Eating Disorder, loved ones, as well as treatment providers and researchers, and to advance the field of Eating Disorders in Canada.

#### Mandate

- We will support the collaborative relationship between the National Initiative for Eating Disorders, the Eating Disorders Association of Canada – Association des Troubles Alimentaires du Canada, the National Eating Disorder Information Centre and the Eating Disorders Foundation of Canada on all matters that pertain to the collective national field of Eating Disorders in Canada.
- We will establish national goals and collaborative activities as determined to be helpful for the advancement of the field of Eating Disorders in Canada.

- **3.** We will determine the level of commitment and financial investment required of each organization prior to the undertaking of any joint activity.
- 4. We will also value the input and participation of provincial and territorial stakeholders as well as other groups of relevance to the field of Eating Disorders.
- 5. We recognize that speaking with one united voice has a stronger impact and will increase the likelihood of successful outcomes. Therefore, we commit to seek out the input of our

partner organizations prior to engaging in national communications or advocacy initiatives focused on Eating Disorders in Canada.

- 6. Collaborative activities could include, but are not limited to, the articulation of a national strategy on Eating Disorders, interorganizational fund-raising activities, other activities of national prominence requiring the support of the field of Eating Disorders.
- **7.** We will evaluate the effectiveness of our interorganizational activities on a regular basis.

## Acknowledgements

The Alliance is grateful for the leadership, time and expertise of the many passionate caregivers, health care professionals, people with lived experience, social service providers and researchers who provided insights and input to reviewing or writing significant sections of this Strategy and its recommendations.

We are also grateful to Dr. Angela Grace, Heather Lilicco, R.H.N. and Dr. Karen Patte who researched and wrote the first version of the Prevention Pillar.

The authors of the strategy, who are representatives or former\* representatives of the Canadian Eating Disorders Alliance, takes full responsibility, though, for the final version of this Strategy and any oversights or errors.

# Canadian Strategy Authors

Eating Disorders Association of Canada-Association des Troubles Alimentaires du Canada (EDAC-ATAC)

Dr. Cheryl Aubie Dr. Nicole Obeid Dr. Josie Geller Dorita Shemie, M.S.W.

National Initiative for Eating Disorders (NIED) Mark Ferdinand Wendy Preskow Lynne Koss

#### Eating Disorders Foundation of Canada (EDFC) Dr. Robbie Campbell Angela Rinaldis

Dr. David Pilon

National Eating Disorder Information Centre (NEDIC) Kelsey Johnston Emily Tam

# Vision

That all Canadians affected by EDs and related mental illnesses have equitable access to high quality, evidence-based publicly covered health care and social services to support their recovery – no matter where they live in Canada.

# The Canadian Eating Disorders Strategy

This strategy contains 6 key pillars and 50 recommendations aimed at improving outcomes for people living with and recovering from EDs and related mental illnesses.

Each pillar is co-chaired by respected researchers, clinicians, caregivers, health care professionals and social service providers from across Canada who conduct research on EDs and related illness, and/or treat and support people directly affected by EDs.

We believe that we can successfully implement many recommendations in this Strategy over the next 10 years

<sup>5</sup> Mental Health Commission of Canada; CAMIMH members; psychiatrists, physicians, psychologists, registered dietitians, social workers, psychotherapists/counsellors, dentists, occupational therapists, physical therapists, nutritionists, naturopaths, acupuncturists, chiropractors, massage therapists, dentists, osteopaths, and other health care professionals; family caregivers including parents and siblings; employers, researchers, teachers, first responders and other professionals.



by adopting a highly collaborative approach involving multiple stakeholders, including people with lived experience, caregivers, health and social services providers, researchers<sup>xiii</sup>, existing health care, mental health and social services organizations, policymakers, governments and other stakeholders<sup>5</sup>. The successful implementation of several recommendations will require, among other activities, the creative exploration, modelling and selection of highly collaborative

approaches to working across different stakeholder groups regardless of service and professional boundaries<sup>xiv</sup>.

# Prevention

# OBJECTIVE

**1.1** To reduce the prevalence and severity of EDs in Canada.

### Introduction

Prevention efforts need to target the time period before someone develops an ED to help divert developing cases and to reduce the severity of a disorder if it does develop.

### **Canadian Context**

Programs and messaging for ED prevention have been developed in the past, but information on effective resources/programs is scattered and research evaluating the effectiveness remains limited. Limited funding is available to conduct research to fill evidence gaps regarding which

targeting other health-related conditions (i.e., obesity) are often given priority over ED prevention, yet anti-obesity messaging has been implicated in the unintended development of ED symptoms<sup>xv</sup>.

activism).

# **1. Prevention Recommendations**

Consistent with Research recommendations made in this Strategy, a well-connected network to disseminate best practices to practitioners and the public is recommended given that Canada is geographically large. Where to focus

prevention efforts (e.g., certain age or demographic groups) or when a tailored approach is needed (i.e. one size fits all messaging will not always work) must also be determined through a robust, ongoing research strategy.

types of programs or what core components

Body Project, feminist-informed approaches,

social justice, media literacy programs, media

Moreover, relatively few researchers within

Canada are focused on ED prevention given

the lack of funding support. Many initiatives

are most effective for prevention (e.g., The

### **PREVENTION RECOMMENDATIONS**

- EDs and a culture influenced by weight bias<sup>xvi</sup>.
- 2. Develop messaging for the general population about prevention.
- 3. Educate practitioners, educators, policy makers, medical and obesity professionals, professionals; dentists; media) on
  - related mental illness:
  - best practices for prevention;
  - appropriate messaging in specific settings (e.g. schools); and
- 4. Identify gaps where no best practices exist, either for certain populations or disorder types.
- in other countries.
- repository to share best practices for prevention-related materials.
- practices.
- 8. Consistent with R. 43, systematically measure efforts aimed at reducing prevalence of EDs in Canada.

<sup>6</sup>This is important, as currently sometimes people with EDs are referred erroneously to obesity clinics, which cause delays in treatment and possibly unintended consequences on the individual being referred for treatment.

1. Develop a common language or definition of what prevention means in the context of

and other relevant groups (e.g., fitness professionals; dance coaches; education

· early identification and identification of people at risk of developing EDs or

timely referral to either information resources or appropriate care<sup>6</sup>.

5. Connect with researchers to fill evidence gaps, including looking to "success" stories

6. Validate whether to use existing best practices repositories or the creation of a new central

7. Subject to R. 6, create a centralized repository to facilitate sharing of better ED prevention

# Public Education and Awareness

# **OBJECTIVES**

**2.1** To reduce stigma and shame associated with EDs; and

**2.2** To better support ongoing awareness and education about EDs in Canada.

## Introduction

Educational interventions targeting stigma and mental health literacy are needed in Canada to change misconceptions about EDs<sup>xvii</sup> held by the public as well as health professionals working with those who may be affected by EDs.

According to a 2015 public survey conducted by Ipsos Reid<sup>xviii</sup> (on behalf of NEDIC), "4 in 10 (40%) Canadians mistakenly believe that EDs are a choice while 21% believe they are genetic".

Moreover, a survey conducted by NEDIC in 2018<sup>7</sup> to gauge literacy about EDs in Canada revealed that among the professionals who participated, the group categorized as "health care providers with high levels of engagement with medical service users (general practitioners, nurse practitioners, and nurses)" did not score statistically significantly higher in ED literacy than the average for professionals as a whole. Also of

note, fitness professionals scored lowest in ED literacy and highest in endorsement of stigmatizing beliefs, yet were most likely to agree that public education about EDs is "adequate". This is concerning, as we know that sociocultural factors, including weight stigma and diet culture, can significantly impact public health as they contribute to risk factors for the development of EDs.

### **Canadian Context**

In Canada, a lack of education as well as misinformation about EDs are barriers to treatment and recovery from EDs. Moreover, low mental health literacy contributes to stigmatization of those with EDs<sup>xix</sup> and may also impede early detection, help-seeking, and appropriate treatment. However, public education can mitigate stigmatization and the barriers to treatment and detection.

<sup>7</sup> NEDIC conducted an online survey of 1000 respondents across Canada to explore the Canadian public's experience with ED education, as well as to explore ED literacy across individuals employed in different professions.



Resources to support additional public education about EDs are scarce, and yet the need is great. A 2012 survey of Canadian teachers<sup>xx</sup> revealed a need for increased provision of education in schools to help teachers to address mental heal challenges in students, including EDs:

- 38.8% of respondents strongly or somewhat agreed that EDs are a pressing concern relatin to student mental health
- Yet 87% of agreed that a lack of adequate staff training in dealing with children's mental illnes is a potential barrier to providing appropriate services for students in their school

While there have been efforts to better educate public about EDs, a strategy including clear guid lines and standards to inform such endeavours does not yet exist. As such, certain approaches that have been identified as harmful for the targ populations may be used in educational efforts

on Ith	individuals or groups who are well-intentioned but not well-informed. Research suggests that the inclusion of direct information and media reports within ED education for young people contra- venes the principle of "first do no harm" xi
ng ff ess	<ul> <li>Providing specific details about EDs results in increased knowledge, but may at the same time result in increased levels of dietary restraint and disordered eating behaviours<sup>xxii</sup></li> <li>The use of media articles (e.g. those in magazines) may also serve as a 'how-to' or information source to adopt disordered eating behaviours due to the perceived glamorization of the disorders<sup>xxiii</sup></li> </ul>
the	
le-	Consequently, there is a need for a significant improvement in the access to and delivery of public education relating to EDs to support early
let by	identification, early access to treatment, and stigma reduction.

# 2. Public Education and Awareness Recommendations

Improved public education related to EDs can facilitate early identification and early access to appropriate treatment, which are actions associated with improved outcomes. Conversely, a

lack of education and misinformation about EDs can create significant barriers for treatment seeking individuals and caregivers who are affected by EDs.

#### **PUBLIC EDUCATION AND AWARENESS RECOMMENDATIONS**

- 9. Seek champions in each province and territory to identify and summarize public education campaigns and best practices. This information will be used to inform the development of evidence-based educational and training materials.
- 10. Using findings from the 2018 NEDIC survey and academic research, further identify significant gaps in knowledge and effective educational programming that could be used to address these specific areas of need.
- 11. Develop and disseminate educational programming to fill gaps in knowledge for appropriate audiences. These tools could include, but are not limited to, webinars, pamphlets, PowerPoint<sup>®</sup> presentations, train-the-trainer resource kits, conferences, and panels. Materials are to be housed in a central location (NEDIC) to ensure streamlined access.

Treatment

# **OBJECTIVES**

**3.1** To improve access to appropriate care in Canada;

**3.2** To reduce barriers to care across the country;

**3.3** 3 To reduce variation in continuum of care for EDs across Canada; and

**3.4** To support outcomes evaluation of care and the development of indicators of recovery.

#### Introduction

ED treatment in Canada is focused on medical stabilization, nutritional rehabilitation, normalization of eating behaviours, and in some cases stabilization of psychiatric comorbidities. Appropriate assessment is required to identify treatment options.

### **Canadian Context**

In Canada, there exist:

• a limited number of specialized, publicly funded EDs clinics or treatment centres<sup>8</sup>, mainly concentrated in urban areas in the

l'Université Laval; and the Nova Scotia Health Authority Eating Disorders Clinic.

(ANEB) Québec; Eating Disorders Nova Scotia; Eating Disorders Foundation of Newfoundland and Labrador; and F.E.A.S.T.

provinces of British Columbia, Alberta, Ontario, Quebec and Nova Scotia;

- Provincial, community and virtual ED networks in British Columbia, Alberta, Manitoba, Ontario, Quebec, Nova Scotia and Newfoundland to help guide people recovering from an ED through the system<sup>9</sup>;
- private clinics and practitioners in most provinces; and
- · little to no services in the Territories.

In Canada's publicly funded health care system, there are varying levels of care available across

<sup>&</sup>lt;sup>8</sup> See, for example, BC Children's Hospital Provincial Specialized Eating Disorders Program; the Provincial Adult Tertiary and Specialized Eating Disorders Program at St. Paul's Hospital; BridgePoint Center in Saskatchewan; the Ontario Community Outreach Program for Eating Disorders (OCOPED), which lists publicly funded programs across Ontario, including the UHN Eating Disorder Program; the Eating Disorder Program at the Douglas Mental Health University Institute; Le Programme des troubles des conduites alimentaires for adults at the Centre hospitalier de

<sup>&</sup>lt;sup>9</sup> See, for example, Kelty Mental Health Resource Centre (BC); The Looking Glass Foundation (Vancouver); the Eating Disorder Support. Network of Alberta; Eating Disorders Manitoba; Sheena's Place (Toronto); Hopewell (Ottawa); Body Brave (Hamilton); Anorexie et Boulimie

the country for people with EDs. Care provided is based on values that are patient- and familycentered, collaborative, interprofessional, evidence-based, and in support of a scientistpractitioner model. However, even when treatment is available, some individuals are not motivated to participate in treatment, which can lead to delays in accessing appropriate care.

In addition to services available in publicly funded organizations, there are several types of specialized services for EDs in Canada, including the following:

- **Residential**: Only a small handful of publicly funded residential facilities exist across Canada. Facilities in Ontario and British Columbia (BC) are intended for those with severe and unrelenting EDs who have tried other treatment options with little to no success. These facilities generally provide services that are short-term in nature, including meal support and around-the-clock care for individuals who are medically stable but who require intensive levels of service to try to combat the ED. BridgePoint Center for Eating Disorders in Saskatchewan also provides publicly funded residential care. However, it is different from the residential centres in Ontario and BC in that it is open to all clients with EDs, not limited to individuals who have achieved little or no success with previous treatment.
- **Inpatient units**: These types of services are available in tertiary care facilities, and house those with an ED who are medically unstable, extremely underweight and those who require intensive support to interrupt their symptoms and/or initiate weight restoration. Inpatient beds can be housed on specialized ED units or are embedded within general medical

units. The care received while an inpatient on a unit involves medical support, nutritional rehabilitation, psychological and psychiatric support and 24-hour care.

- Day Treatment Programs: Also housed in tertiary care facilities, day treatment programs, also known as partial hospitalization programs, are services for those who are medically stable but who require intensive levels of support in order to try to recover from an ED. These intensive programs require individuals to commit to participating in treatment for 6-10 hours every day for 4-5 days per week, and constitute meal support, nutritional rehabilitation, psychological treatment, and addressing comorbid psychological issues. For those who initiated their treatment on an inpatient unit, these programs are often offered and serve as a step down from intensive levels of care. These are group-based programs that usually have 6-10 individuals in care simultaneously.
- Intensive Outpatient Programs (IOPs): These programs are relatively new to Canada. They are generally offered outside of hospitals and may involve visits with different providers or professionals (in a centralized location or different locations). IOPs are intensive programs that provide multiple hours of treatment in a given week. IOPs aim to support those who require less intensive treatment than a day treatment program but more services than one treatment contact per week. These services usually involve one psychological group-based service and one meal provided during each interaction.
- Outpatient programs: These programs tend to be housed in hospitals and provide group-



based and individual therapy. Treatment is provided by an interdisciplinary team and include group-based psychological treatments, and medical, nutritional and psychiatric monitoring

- Step-down outpatient services: These programs that are housed in hospital are a stepdown service for inpatient and day treatment programs and offer individuals on the path to recovery further treatment in the form individ ual psychological therapy and medical/dieteti monitoring at a frequency of once/week.
- **Transitional/Supportive Housing**: There are very few transitional housing opportunities in Canada where individuals, who have completed more intensive training, reside until they



<b>D</b> -	are able to return to their home communities.
es	Sometimes these services are adjacent to more specialized ED programs that continue to pro-
ıg.	vide treatment allowing patients a supportive living environment to further their recovery.
-	
	General, pediatric and adult ED programs dif-
t	fer not only in location, availability of appropri-
)	ate interprofessional teams, and services offered,
-k	but also in treatment philosophies, modalities
ic	used, the involvement (or not) of families, and
	treatment inclusion and exclusion criteria.
e	This fragmented system results in individuals

Chronic ED patients often have complex health histories. Many of the community resources cannot accommodate complex cases. In fact, these are excluded from many of their mandates. In our experience, only the fully implemented inter-disciplinary ED specialty teams were effective.

patients fall through the cracks entirely because their co-morbid symptoms exclude them from admission or their age at presentation of treatment does not match the available services in their geographical region.

Regarding age, **transition from child/youth to adult services** is also challenging and/or disconnected. It is known that treatment transitions within the system are disjointed and often difficult for the patient and their caregivers to navigate, let alone access, or even know about.

There are also an alarmingly high number of **under-resourced populations** (e.g. rural, LGBTQ people, males, indigenous and Northern communities, cultural and ethnic minorities).

Moreover, some treatment facilities do not treat Binge Eating Disorder (BED). In some cases, patients are referred by some primary care providers to a weight management clinic, which focuses on weight loss, and not treating the actual ED, a situation which may further perpetuate the problems associated with BED.

Finally, there is a lack of programming to address commonly occurring comorbidity issues for people with an ED. This in turn results in treatment provided in silos and patients having to move back-and-forth between service providers, leading to ineffective and disjointed care. "Chronic ED patients often have complex health histories. Many of the community resources cannot accommodate complex cases. In fact, these are excluded from many of their mandates. In our experience, only the fully implemented inter-disciplinary ED specialty teams were effective."<sup>10</sup>

Sometimes, individuals with an ED seek services away from their hometown, and sometimes in other countries, notably the United States.

### Therapeutic Approaches and Treatments

There are a number of different treatments that have been proposed for individuals with EDs.

Early treatment of EDs, such as rapid early intervention programs, has been shown to lead to better recovery outcomes, particularly in the length of illness and likelihood of recovery<sup>xxiv</sup>. Similarly, professional educational interventions have been shown to positively change the attitudes of clinicians towards specific EDs therapy<sup>xxv</sup>.

Unfortunately, some approaches to treatment have mixed or limited evidence of effectiveness related to long-term outcomes<sup>xxvi</sup>. However, the evidence base available for **Enhanced Cogni***tive Behaviour Therapy* (CBT-E) – used to treat



adults – and *Family-Based Treatment* (FBT) – used to treat adolescents – is strong. In addition to pharmacotherapy (iii), CBT-E and FBT are the two most commonly used approaches in treatment centres across Canada.

#### i. **CBT-E** or **Enhanced Cognitive Behaviour**

**Therapy**: This approach is concerned with the processes that maintain ED psychopathology rather than those responsible for its initial development. Therefore, the focus is on maintaining mechanisms which help construct a formulation which is then used to target the features that need to be addressed in treatment. CBT-E is generally of fixed length i.e. 20 sessions over 20 weeks. It has 4 well defined stages: Stage 1 intensive initial stage, appointments are twice per week and

<sup>10</sup> Caregiver, NIED 2018 Caregiver Needs Assessment Survey (2018).

it aims to engage the patient in treatment and change. Stage 2 is a transitional stage where we take stock, review and revise formulation. Stage 3 addresses main mechanisms that are maintaining the patient's ED. Stage 4 focuses on the future and minimizes the risk of relapse in the long term.

- ii. **FBT** or *Family-Based Treatment*: Maudsley FBT originated at the Maudsley Hospital in England. It involves intensive outpatient treatment in which parents play an active, positive role in their child's recovery from an ED.
- In addition, there are three guidelines, further developed by Dr. James Lock *et al.*, specific to treating a restrictive ED:

Treatment

- Phase 1: Parents are taught strategies to gain control over the youth's eating behaviours.
- Phase 2: Family issues are discussed and addressed.
- Phase 3: Getting the patient's healthy development back on track.
- FBT further stipulates that
  - the family take a compassionate stance toward the illness;
- the clinician
  - raises anxiety about the illness in the parents as a means of inciting them to take action;
  - lifts blame and guilt that anyone caused the EDs;
  - empowers parents to be an active agent in recovery; externalizes the illness; and
  - places importance on the psycho-education of the illness for the whole family.

iii. **Psychopharmacology**: There is no specific medication meant to treat EDs directly. However, pharmacology can be used to address complex comorbid conditions and symptoms.

#### **Complex Comorbidities**

Common comorbidities of mood, anxiety, substance use, post-traumatic stress and personality disorders are often seen concurrently in those with an ED. Depression or mood symptoms have been found to be the most prevalent comorbidity in those diagnosed with anorexia nervosa (AN) or bulimia nervosa (BN), although recent estimates suggest that various comorbidities exist with EDs with alarmingly high rates: 93% of those with an ED will also experience a comorbid mood disorder; 56% will experience a concurrent anxiety disorder; and 22% will present with a comorbid substance use disorder<sup>xxvii</sup>. As a result of these complex comorbidities, treatments must target not only the symptoms associated with the ED, but also these other mental illnesses.

# **3. Treatment Recommendations**

Canada has various, although limited, ED treatment facilities providing treatment as best as possible for individuals and families affected by an ED. Unfortunately, access to care is not equal across the country, and there are some provinces that have no treatment centres available

for those with EDs. Emergency departments are sometimes used as a first point of entry into the healthcare system for many "and are considered a 'safety net' in response to the lack of other psychiatric services"xxviii.

# **TREATMENT RECOMMENDATIONS**

- intensity and for patients of all illness severities;
- (e.g. Short Treatment Allocation Tool for EDs (STATED))xxix.
- professionals, providers / health care and social services.
- practices.
- and collaborative care.
- 19. Establish a Task Force to improve transitions from youth to adult services.
- identified early.
- sub-populations.
- (e.g. obesity clinics).
- 24. Develop and deliver specialized training to optimize treatment planning for and the integration of treatment(s) that addresses complex comorbidities.

12. Consistent with principles in the Canada Health Act, publicly state that all Canadians, including individuals living with an ED, must have equitable access to medically necessary services to all Canadians – no matter where they live and without discrimination.

Treatmen

13. Create, fund and implement specialized services to address EDs across all levels of treatment

14. Support spread of leading practices aimed at better matching of treatment to patient needs

15. Improve communication and sharing of best practices and evidence-based care across

16. Design networked services for EDs across Canada based on promising and evidence-based

17. Establish networks in provinces and territories where they do not exist to coordinate resources

18. Establish a pan-Canadian 'network of networks' to link together these networks across provinces and territories to further support dissemination of best practice and coordination of care.

20. Develop stronger partnerships with primary care to ensure that individuals with EDs are

21. Develop bilingual, culturally appropriate and gender-sensitive<sup>xxx</sup> services for marginalized

22. Develop and deliver training in sensitivities regarding EDs in non-ED treatment environments

23. Encourage provincial health authorities to support participation of Eating Disorder clinicians in learning opportunities to enhance their treatment competencies (e.g. EDAC-ATAC conference).

# **Caregiver Support**

# **OBJECTIVES**

**4.1** To connect caregivers with evidence-based and evidence-informed resources and information to

- help them better navigate health care system and social services;
- support their own health;
- support the recovery of the person they are caring for; and

**4.2** To develop and propose improvements to existing health, workplace and social policies and programs so that they are more responsive to caregiver needs.

# Introduction

Caregivers, along with the person they are caring for, along with various professionals, providers and other family members, play an important role in supporting the treatment of a person diagnosed with an ED. This is consistent with clinical practice guidelines for EDs<sup>xxxi</sup>, which promote team-based approaches to care.

# **Canadian Context**

Informal caregivers are unpaid caregivers. They play an important part in improving the wellbeing of people living with a mental illness. According to the Mental Health Commission of Canada (MHCC), "Caregivers often provide a variety of supports, including navigating the mental health system to access services, transportation, advocating for services, and social, financial, and emotional supports. Studies have shown that caregivers can experience a significant sense of personal growth, fulfillment, and purpose through the support they provide to a friend or family member with a mental health problem or mental illness. At the same time, the caregiver's own health — both physical and emotional can be negatively affected by the day-to-day demands of long-term caregiving."xxxii



**C** Caregivers often provide a variety of supports, including navigating the mental health system to access services, transportation, advocating for services, and social, financial, and emotional supports. Studies have shown that caregivers can experience a significant sense of personal growth, fulfillment, and purpose through the support they provide to a friend or family member with a mental health problem or mental illness. **9** 

However, this role is only starting to be understood by society. The MHCC acknowledges that "more needs to be done to support caregivers, both by acknowledging and addressing their needs, and reducing the stigma they still experience".xxxiii

Although there are no comprehensive data sets research in Canada regarding the role of caregivers ers of people affected by EDs<sup>xxxiv</sup>, we know that caregivers of people living with a mental illness want to play a pivotal role in the recovery of the person they are caring for<sup>xxxv</sup>. We also know that multiple stakeholders, including caregivers of people living with a mental illness, have recently developed guidelines and standards to improve mental health services in Canada.

od	Caregivers of people living with an ED expe- rience, as many Canadians with complex care
	needs, wait for appropriate care and do their
by	best to cope while waiting: "The local children's
	hospital ED program currently has a 4 week wait
	list. Our family doctor is monitoring our daugh-
s or	ter's vitals on a weekly basis but is unsure of
V-	what kind of advice to give. If the ED program
	or someone could provide a "so your child has
5	an ED, and you're waiting for treatment - here's
5	what to do", that would be so helpful. I am
t	cobbling together a plan from online forums and
	published articles, and hoping it works ok."xxxvi
ly	
e	To help inform this Strategy, NIED surveyed care-
	givers of people living with or recovering from

EDs in 2018<sup>11</sup> to better understand their unmet needs as well as their successes in caring for themselves while they cared for someone living with an ED. Consistent with findings from various Statistics Canada surveys<sup>xxxvii</sup> on caregiving, caregivers who responded to this survey were more likely to be anxious, constantly worried, stressed, exhausted and/or depressed while caring for their loved one. Other caregivers experienced isolation, helplessness, poor sleep, mood swings, self-doubt and loss of autonomy as a result of caring for some affected their family member recover from an ED. For by an Eating Disorder.

Understanding the relationship between the impact of caregiving on a caregiver's health status<sup>xxxviii</sup> is important, as the potential negative effects include an inability to care for their loved one, weaker attachment to the workforce, and increased use of health care services.

"Findings relating increased family burden to increased presence of personal mental health difficulties conform to and extend existing research among caregivers. This further stresses

the need for the investment of resources in the protection of the well-being of this group who perceive family burden, for moral and economic reasons or practical reasons, such as the impact of their mental health on their continued provision of care and the cared for person's outcome"xxxix.

Just over half of caregivers responded that they understood what they needed to do to support those caregivers who were employed, about onethird (1/3) of caregivers said they were" very satisfied" or "satisfied" with the quality of support or care they receive from or through their employer.

Finally, caregivers provided a number of suggestions on how care and supports can be improved, including increased funding, a greater focus on the person requiring care, increased access to resources and financial assistance to help with daily living, better education for professionals and access to family or parent support groups, especially in rural and remote locations.

# 4. Caregiver Support Recommendations

The following recommendations are consistent with recommendations in the MHCC's National Guidelines for a Comprehensive Service System to Support Family Caregivers of Adults with Mental Health Problems and Illnesses<sup>xl</sup> and observations shared by caregivers with NIED and in various Statistics Canada surveys over the last 10 years. "The Mental Health Strategy for Canada recommends more help for family caregivers through better financial supports, increased access to respite care and more flexible workplaces. Planning a comprehensive approach to meeting the needs of family caregivers is not the sole responsibility of the mental health system and requires action on the part of many other system stakeholders, including governments and other organizations."xli

<sup>11</sup> Qualitative responses to an online survey were received from 67 caregivers living in 9 out of 10 provinces.

**C** The Mental Health Strategy for Canada recommends more help for family caregivers workplaces. Planning a comprehensive approach to meeting the needs of family caregivers is not the sole responsibility of the mental health system. **))** 

### **CAREGIVER SUPPORT RECOMMENDATIONS**

- Strategy.
- when appropriate, as part of the care team.
- Living with Mental Illness or Experiencing Mental Health Challengesxiii.
- absences from work due to a need to fulfill caregiver responsibilities.
- support of caregivers of people living with mental illness.



through better financial supports, increased access to respite care and more flexible

25. Engage caregivers formally in the design and development of recommendations in this

26. Validate and prioritize with caregivers and other stakeholders how this Strategy can support the implementation of recommendations in the National Guidelines for a Comprehensive Service System to Support Family Caregivers of Adults with Mental Health Problems and Illnesses.

27. Continue to work collaboratively with the MHCC's Family Caregiver Advisory Committee and national and provincial caregiver organizations to support efforts aimed at improving caregivers' access to resources and information about mental illness, involvement in system and program planning; navigating health care systems and social services; and engagement,

28. Develop, with national and provincial caregiver organizations, new or adapted provincial and organizational policies and programs that will provide better support to all caregivers who need respite and/or who are trying to access appropriate care for their loved ones.

29. Further examine with caregivers and stakeholders the value and effectiveness of existing caregiver guides such as the Mental Health Caregiver Guide: A Guide for Caregivers of Persons

30. Develop new or adapt existing workplace policies that can better accommodate periodic

31. Work with employers to identify better practices that can be shared, spread and adapted in

# **Training**

# **OBJECTIVES**

5.1 To identify post graduate and continuing professional development training pathways for EDs;

**5.2** To develop clinical competencies related to the provision of health services to individuals with EDs and to their families; and

**5.3** To encourage future graduates to pursue research careers in the area of FDs.

# Introduction

5

Best practice ED treatment is interprofessional<sup>xliii</sup>. Medicine (Family Practice, Pediatrics [including Adolescent Medicine], Psychiatry), Psychology, Dietetics, Nursing, Advanced Practice Nursing, Occupational Therapy, and Social Work disciplines work closely with people, their caregivers and their family members to contribute to ED care. Other disciplines including Child and Youth Workers, Art Therapists, and Teachers may also work in ED care settings.

Professional training is integral to supporting advanced learning and practice related to the provision of high-quality care for EDs. Training can be used to support learning in several areas, including, but not limited to: research, early identification, prevention, health promotion,

assessment, referrals, diagnosis, care planning and coordination, treatment selection, medication management, counselling, surveillance and more.

Through training, practitioners can acquire both foundational skills in post-graduate settings related to the provision of high quality, professional health care service delivery and ED specific competencies which are generally not available as part of graduate curricula across Canada.

## **Canadian Context**

In addition to limited graduate course offerings specialized in EDs, Canadian professionals receive or acquire specialized skills regarding EDs from a variety of people and sources:



- Health care professionals who provide services to individuals with EDs are typically oriented, mentored and/or supervised by more seasoned providers working in those services.
- Service settings may also support these providers to seek continuing professional education to enhance ED knowledge and skills through a variety of on-the-job or intensive training experiences. These could include:
  - workshops and online learning modules offered by ED specialists or provincial discipline-based associations;
  - National/international conferences such as the Eating Disorders Association of Canada – Association des Troubles Alimentaires du Canada. the International Conference on

Eating Disorders (ICED), the London Conference on Eating Disorders;

- opportunities to job shadow another professional in specialized ED programs;
- intensive training offered through supervised opportunities.

## **International Training**

Canadian ED professionals may also have membership in the International Association of Eating Disorder Professionals (iaedp<sup>™</sup>), which provides certification of ED professionals to promote standards of excellence in the field of EDs. Professionals must demonstrate clinical expertise through education, experience and an exam to earn a certificate in a variety of discipline-based categories.

# **5. Training Recommendations**

Despite having some clinical guidelines and evidence-based treatments available to guide practice, there is insufficient training in evidence-based treatments available across Canada

and very few lead Canadian trainers in various treatments for mentoring purposes. This leads to few practitioners across the country having access to training in evidence-based practices.

#### **TRAINING RECOMMENDATIONS**

- 32. Coordinate and disseminate training in evidence-based treatments as they become available for current professionals.
- 33. Develop a continuing professional development webinar series by Canadian ED specialists with appropriately recognized learning credits.
- 34. Deliver training in sensitivities regarding EDs in non-ED treatment environments, which could include correctional settings and addictions.
- 35. Disseminate best practices via Telehealth for rural and remote areas.
- 36. Enhance training in Family Practice settings, primary care networks/health teams and integrated health networks to support ED screening and early identification.
- 37. Support professionals wishing to transition their practice into the field of EDs via intensive exposure within specialized ED settings.
- 38. Identify training/exposure opportunities to other healthcare professionals who come into contact with individuals with EDs (e.g., Gastroenterologists, Emergency Department staff, Paramedics, etc.).
- 39. Create a task force to identify minimum standards for graduate education in EDs within all health disciplines (including development of a standard curriculum for EDs).
- 39. Create a task force to identify minimum standards for graduate education in EDs within all health disciplines (including development of a standard curriculum for EDs).

# Research

# **OBJECTIVES**

**6.1** To identify research priorities, needs and gaps in the field of EDs in Canada;

**6.2** To enhance Canadian collaborative research and support for EDs research in Canada; and

**6.3** To enhance ongoing surveillance, performance measures and data related to EDs.

#### Introduction

Ongoing research in the field of EDs is vital to better understand the etiology of EDs, as well as prevention and evidence-based treatment approaches. It is equally essential to effectively translate research findings to improve care and outcomes for those affected by EDs.

#### **Canadian Context**

A strength of the Canadian research landscape is the publicly funded healthcare system allowing for treatment to be determined by best practices and clinical guidelines as opposed to other factors. This should allow for the promotion and implementation of clinical research to inform practice. Accordingly, Canada has several clinician-scientists across the country who are promoting EDs research, and some provincial gover ments have invested meaningfully and support research related initiatives in the field of EDs<sup>12</sup>.

<sup>12</sup> For example, see notably, Ontario's investments in the Ontario Community Outreach Program for Eating Disorders (OCOPED) since 1994, http://www. ocoped.ca/ and the establishment of an integrated approach to addressing EDs in British Columbia, started in 2009, which resulted in the adoption of an Action Plan for Provincial Services for People with Eating Disorders, the adoption of Clinical Practice Guidelines for the BC Eating Disorders Continuum of Services, the creation of a Centre for Excellence for Eating Disorders and the expansion of the BC Eating Disorders Community of Practice (CoP).

	Unfortunately, the field of EDs research continues
	to be vastly under-funded, and consequently
	under-studied. Of the limited funding that
	exists in Canada, it is usually not available for
	the field of EDs. A 2014 reportli of the Standing
	Committee on the Status of Women suggests
	that across a 5-year period, \$7.5 million has been
	spent on operating grants for EDs in Canada,
	while \$86 million has been spent on operating
	grants for schizophrenia, despite EDs being as
S	severe and more prevalent and deadly than
	schizophrenia. This same report also states that
	students, trainees and early career scientists are
	being coached to stay away from specializing
	in EDs to avoid career pitfalls resulting from
	this marginalized area of health. Practically, this
	means that there are few research mentors/
	supervisors/investigators available. There
rn-	are many students across various disciplines
ed	interested in studying EDs, but there are few
	research training opportunities and few graduate





program research supervisors able to support advanced research training across the country, limiting the emergence of new Canadian scientists in the field and advancement and innovation in the field.

With regard to research translation, clinical research findings are not reaching clinicians across the country, preventing individuals with EDs from benefiting from these areas of strength and innovation. There are no dedicated cross-country research bodies that exist in the field of EDs (e.g. peer-reviewed journals, clinical information service such as UpToDate<sup>xliv</sup>). Such a network is essential to engage in pan-Canadian

research as well as with various cultural and indigenous communities to enhance collaboration, research possibilities and knowledge translation that can inform practice and improve outcomes.

Finally, in Canada, as elsewhere in the world, statistics related to EDs in primary care are generally regarded as poor<sup>xiv</sup> or woefully out-of-date.

On July 3, 2018, federal, provincial and territorial (FPT) health ministers endorsed CIHI's plans to "measure pan-Canadian progress toward improving access to mental health and addictions services and to home and community care"xivi. Provincial and territorial governments' development of new mental health and community care indicators<sup>13</sup> represents a great opportunity for stakeholders to identify relevant or new data sources that can support comparable reporting of performance across Canadian jurisdictions.

The creation of these new methods of data collection and performance measurement will

# 6. Research Recommendations

There is strong potential for research advancecarers and clinicians. This project provides the ment in EDs from Canadian researchers, as field with a consensus-built understanding of strong collaborative research teams are emergwhere research efforts should be placed next. ing across the country and Canadian research is Canadian researchers are in a unique position to well-represented at international conferences. leverage this work and to advocate to research Greater investment could accelerate this expofunders for more investment in these priorities. nentially. However, there are no research chairs in Canada for EDs research to increase the It is relevant to broaden the scope of ED research, as it is fairly homogenous, in that research capacity of Canadian research. In 2018, a panexists mostly for women with anorexia nervosa, Canadian patient-oriented research project<sup>xlvii</sup>, funded by the Canadian Institutes of Health whereas research in other diagnostic categories Research (CIHR), identified the top 10 research or in differing populations (e.g. adolescents, priorities from those with lived experience, their males) remains limited.

### **RESEARCH RECOMMENDATIONS**

40. Continue to identify research needs and priorities by people with lived experience,

<sup>13</sup> Except Quebec and Ontario. "The federal government has agreed to an asymmetrical arrangement with Quebec, distinct from the Common Statement of Principles. In addition, given the transition to a new government in Ontario, the province cannot officially endorse the recommendations," https://www.newswire.ca/news-releases/a-canadian-first-cihi-to-measure-access-to-mental-health-and-addictions-services-and-to-home-and-community-care-687245651.htm

lead to improved knowledge exchange related to mental illnesses, including EDs. The use of comparable, pan-Canadian data sets and measures will also greatly enhance the potential for jurisdictions to learn from each other's successes and challenges and continuously improve the responsiveness and accessibility of existing mental health resources aimed at preventing and treating complex mental illnesses.

Researc

caregivers, and health care professionals in the field so that research questions of most importance to those affected by an ED are carried out and widely promote these priorities<sup>XIII</sup>.

### **RESEARCH RECOMMENDATIONS** (cont'd)

- 41. Support a systematic approach to the identification of research priorities across the spectrum of EDs and affected populations.
- 42. Update<sup>xlviii</sup> existing EDs data and relevant indicators in Canada.
- 43. Update existing mental health performance measures<sup>xiix</sup> so that they include EDs<sup>1</sup>.
- 44. Enhance uptake of available ED research by using various knowledge translation strategies and models.
- 45. Identify/create research training opportunities across all disciplines.
- 46. Encourage collaboration, organization and planning of research objectives and combination of expertise to attain federal (i.e., CIHR, SSHRC) funding to carry out pan-Canadian research.
- 47. Establish a Canadian Research Chair for EDs.
- 48. Establish a Canadian Research Network for EDs, modelled on the EDs Centres for Excellence in British Columbia.
- 49. Increase financial support for research related to EDs.
- 50. Create a national data repository for enhanced research capabilities that reflect the Canadian context.

# Conclusion

(Registered Psychotherapist), NIED brought **How this Strategy Was Developed** together approximately 40 stakeholders from In 2015, following the publication of a thoughtful 9 out of 10 provinces (ED clinicians, researchers, report entitled<sup>II</sup> Eating Disorders among Girls and frontline health care providers, parents with *Women in Canada* by the House of Commons lived experience, caregivers and volunteers). Standing Committee on the Status of Women in Stakeholders participated actively in a workshop 2014, EDAC-ATAC, EDFC, NEDIC, and NIED14 disto define a focus for a pan-Canadian EDs stratcussed how we could collaborate meaningfully egy. Many volunteered to further refine possible to improve outcomes for people living with actions we could take collectively to improve EDs in Canada. outcomes for people affected by EDs in Canada. We would like to sincerely thank them for their insights, expertise and encouragement, which These conversations led to the development of a survey to systematically engage stakeholders have inspired us throughout this journey. across Canada to identify gaps that could be worked on to better support care and recovery Following the conference, four national ED for individuals and others affected by EDs. organizations — EDAC-ATAC, EDFC, NEDIC and NIED — agreed to work together to create a Prior to the September 2016 EDAC-ATAC concomprehensive set of actions that would fill gaps ference, NIED worked with stakeholders and identified by stakeholders who participated in a volunteers, led by Andrea Lamarre, to develop national survey in 2016. and disseminate a survey to approximately 500 private and public organizations across Canada,

These actions appear as 'Recommendations' grouped under 6 pillars) throughout this Stratincluding hospital programs, private practices, mental health organizations, and all those who egy. We believe these recommendations will participated in NIED's M.O.M. (Mothers & Others) improve outcomes for people living with or March Against EDs on October 27<sup>th</sup>, 2015. Based recovering from EDs in Canada. on responses from 400 stakeholders, we identi-This first Strategy will be used to expand our dialogue with stakeholders, refine specific recfied 6 pillars: Training, Education, Treatment, Prevention, Advocacy and Research. Following ommendations and develop action plans for the the EDAC-ATAC conference, a 7<sup>th 15</sup> pillar was Strategy's implementation starting in 2019. added: Caregivers.

With the support of conference organizers, Dr. Heather Wheeler and Carly Crawford

<sup>14</sup> Eating Disorders Association of Canada – Association des Troubles Alimentaires du Canada (EDAC-ATAC); National Eating Disorder Information Centre (NEDIC); Eating Disorders Foundation of Canada (EDFC); and National Initiative for Eating Disorders (NIED).

<sup>15</sup> Since that time, 'Advocacy' is now seen as a separate but related step to implementing this Strategy and its recommendations. As such, it is not a pillar in this Strategy. Given that a wide range of stakeholders will be needed to take concrete actions to implement recommendations in this Strategy in jurisdictions across Canada, we cannot predict at this time how implementation, let alone Advocacy, will occur.

We look forward to future dialogue with stakeholders to discuss how best to implement this Strategy's recommendations.

# **Annex 1**

# Why create a Canadian Eating Disorders Strategy?

ince the 1950s, the prevalence of EDs have been increasing around the world, and mainly in western and westernized countries<sup>lii</sup>. Moreover, governments around the world underinvest in EDs and mental health care compared to other services that serve relatively few people<sup>liii</sup>.

Unfortunately, too many Canadians with EDs do not have access to timely, appropriate care, contrary to at least 2 out of the 5 principles of the Canada Health Act - universality and accessibility, which state that all Canadians are entitled to "the same level of health care" and have "reasonable access to health care facilities."

Based on a 2018 survey of caregivers, we know that navigating existing health care system and social service organizations and accessing timely, high quality specialized care remains a challenge for many people. One caregiver's frustration clearly illustrates the challenge they face when trying to find help: "I have been turned away from the hospital on several occasions even after being sent to [the emergency department] at the family doctor's request because her ECG showed that her heart was failing. Just this past winter we were at the hospital four times and turned away all four times, she is 24 and weighed 72lbs. We were terrified we were going to lose her! We finally got her into a private treatment centre and she was sent home because she was "too sick to be there". Too sick for a treatment centre. not sick enough for the hospital.... That is "extremely stressful". When you become solely

responsible for your child's life with absolutely no support from the medical community the pressure is enormous, we are often expected to do what the system can't. The illness is often misunderstood by medical professionals and the resources for care are slim to none. Caring for someone with an eating disorder is an overwhelmingly stressful responsibly. We have been at this for eight years."liv

Suicide and suicidal ideation remain a challenge for many people struggling with an ED or related mental illness, such as depression, anxiety or an addiction. Members of the Alliance are actively working with the pan-Canadian Suicide Prevention Collaborative to reduce the rate of suicide for all Canadians and give life to the Federal Framework for Suicide Prevention Act<sup>IV</sup>, adopted by the Parliament of Canada in 2012.

Regarding education and training, which we believe are foundational to achieving many of the recommendations contained within this Strategy, we aim to work collaboratively with health professionals, providers, caregivers and other stakeholders to design and develop new, evidence-based education and training courses to reduce risks, promote stronger health and speed recovery from EDs.

Based on survey responses of medical doctors<sup>16</sup>, 70% of doctors receive 5 hours or less of ED-specific training while in medical school<sup>Ivi</sup> and only 6.3% of psychiatry residents felt they had spent enough time with patients with EDs to work effectively with them in clinical practice<sup>lvii</sup>.

<sup>&</sup>lt;sup>16</sup> In 2018, NIED worked with the Dietitians of Canada to better understand their educational needs related to the treatment of EDs. From this survey and ongoing engagement with dietitians and other professionals, we aim to develop and launch a similar survey with a wider group of health professionals in 2019 to better understand their needs for university training and continuing professional education (CPE) and continuing medical education (CME). The results of these training needs assessment surveys will complement what we expect to learn from a survey launched in July 2018 aimed at caregivers and non-professionals to better understand their knowledge and general educational needs associated with EDs.

This Strategy aims to update our understanding of learning needs of the general public and several professional disciplines. We aim to develop educational and training offerings that can fill knowledge gaps and improve practice.

Overall, multi-disciplinary research, education and training will help us to:

- reduce stigma associated with mental illness;
- avoid unintended harms associated with certain behaviours or practices in society and care; and
- better share and use evidence to guide specific interventions aimed at preventing or treating EDs and related mental illnesses.

Moreover, there are very few<sup>17</sup> examples of models of coordinated and continuous care for EDs in jurisdictions across Canada or around the world. In Canada, this has resulted in people seeking specialized medical and residential care for EDs outside of Canada, at great social and emotional expense to families and financial expense to Canadian health care systems<sup>IVIII</sup>. We know that collaborative models are important to the timely identification of risk factors and referral to and use of appropriate support or care, all of which aim to prevent chronicity of symptoms, improve quality of life, improve chances of recovery and save lives through treatment.

Finally, the social and economic costs associated with EDs are large. In 2003, a study reported that provincial costs of people living with AN

on long-term disability in British Columbia may be as high at \$101.7 million/year (30 times the cost of all provincial specialized ED services)<sup>lix</sup>. Direct health system costs for treating EDs were estimated at \$99.9 million in 2012 in Australia<sup>lx</sup> and productivity loss for society estimated at \$15.1 billion<sup>lxi</sup>. While similar financial impact analysis is not available in Canada on a national scale, prevalence rates for EDs in Canada mirror those of Australia.

Based on recently published research regarding the cost effectiveness of prevention and treatments for EDs<sup>18</sup>, as well as the value of spending on social services<sup>lxii</sup> in Canada, we are optimistic that we can explore options with stakeholders to better to allocate resources across health and social spending to achieve better outcomes for people affected by EDs and related mental illnesses in Canada.

#### Building on Existing Federal, Provincial and Territorial (FPT) Investments in Mental Health and Community Care

Multiple organizations, governments and individuals are already working together to prioritize research<sup>Ixiii</sup> as well as increase awareness and education about EDs in Canada<sup>Ixiv</sup>.

We believe that we can successfully build on these foundational collaborations by exploring with stakeholders ways to better apply research to practice.



We are equally optimistic that new investments being made by provinces and territories in expanded community mental health programs and home care will have a positive impact on pe ple with EDs and their caregivers, even though i is recognized that EDs require specialized, tertia levels of care, which remain underfunded and under-resourced across Canada.

Over the last two decades, provincial and territorial governments have made important investments in community care, mental health and home care. In late 2016 and early 2017, the Government of Canada signed 10-year funding agreements totalling \$11.5 billion with province and territories to specifically improve home care and mental health initiatives across Canada.

As provinces and territories continue to invest in community mental health and home care initiatives, opportunities exist to strengthen existing



eo-	social support programs for caregivers of people affected by EDs and to support the integration of evidence-informed practices into existing and updated mental health programs.
it ary	Both collaboration and increased engagement with multiple stakeholders will lead to a more efficient and effective use of existing govern- ment resources invested in community mental health and health care across Canada.
es e n	We also believe that highly collaborative approaches to knowledge dissemination and mobilization better and promising practices on existing, well-used knowledge exchange plat- forms (such as the Canadian Best Practices Portal, managed by the Public Health Agency of Canada, the B.C. EDs Community of Practice, the Evidence Exchange Network for Mental Health and Addic-
-	tions, or the Knowledge Exchange Network <sup>Ixv</sup> maintained by Children's Healthcare Canada).

<sup>&</sup>lt;sup>17</sup> In Canada's most populous province, Ontario, the Ontario Community Outreach Program for Eating Disorders (OCOPED) is one example of a provincial network of specialized ED service providers.

<sup>&</sup>lt;sup>18</sup> "Eating disorder interventions were mainly cost-saving or more effective and more costly compared to comparators; however, some results did not reach statistical significance": Khanh-Dao Le, L., Hay, P and Mihalopoulos, C. "A systematic review of cost-effectiveness studies of prevention and treatment for eating disorders," *Australian and New Zealand Journal of Psychiatry*, April 2018, Volume: 52 issue: 4, page(s): 328-338, <u>https://doi.org/10.1177/0004867417739690</u>

# 1. Prevention Recommendations

Objective	Re	commene
<b>1.1</b> To reduce the prevalence and severity of (EDs) in Canada.	1.	Develop a means in t weight bia
	2.	Develop m preventior
	3.	Educate pr and obesit fitness pro dentists; m • early ide developi • best prac • appropri • timely re care <sup>6</sup> .
	4.	ldentify ga populatior
	5.	Connect w looking to
	б.	Validate w the creatio for prevent
	7.	Subject to sharing of
	8.	Consistent reducing p



# Summary of Recommendations

## dations

a common language or definition of what prevention the context of EDs and a culture influenced by as<sup>xvi</sup>.

messaging for the general population about n.

practitioners, educators, policy makers, medical ity professionals, and other relevant groups (e.g., ofessionals; dance coaches; education professionals; media) on

entification and identification of people at risk of bing EDs or related mental illness;

actices for prevention;

riate messaging in specific settings (e.g. schools); and eferral to either information resources or appropriate

aps where no best practices exist, either for certain ons or disorder types.

with researchers to fill evidence gaps, including "success" stories in other countries.

whether to use existing best practices repositories or on of a new central repository to share best practices ntion-related materials.

R. 6, create a centralized repository to facilitate better ED prevention practices.

t with R. 43, systematically measure efforts aimed at prevalence of EDs in Canada.

# 2. Public Education and Awareness Recommendations

Objectives	Recommendations		
<b>2.1</b> To reduce stigma and shame associated with EDs; and	9. Seek champions in each province and territory to identify and summarize public education campaigns and best practices. This information will be used to inform the development of evidence-based educational and training materials.		
<b>2.2</b> To better support ongoing awareness and education about EDs in Canada.	10. Using findings from the 2018 NEDIC survey and academic research, further identify significant gaps in knowledge and effective educational programming that could be used to address these specific areas of need.		
	11. Develop and disseminate educational programming to fill gaps in knowledge for appropriate audiences. These tools could include, but are not limited to, webinars, pamphlets, PowerPoint <sup>®</sup> presentations, train-the-trainer resource kits, conferences, and panels. Materials are to be housed in a central location (NEDIC) to ensure streamlined access.		

# 3. Treatment Recommendations

Objectives	Recommend
<b>3.1</b> To improve access to appropriate care in Canada;	12. Consistent state that a ED, must ha services to
<b>3.2</b> To reduce barriers to care across the country;	without dis
<b>3.3</b> To reduce variation in continuum of care for EDs across Canada; and	13. Create, fund EDs across all illness se
<b>3.4</b> To support outcomes evaluation of care and the development of	14. Support sp of treatmer Tool for EDs
indicators of recovery.	15. Improve co evidence-b care and so
	16. Design net promising a
	17. Establish ne do not exis
	18. Establish a together th to further s coordinatic
	19. Establish a adult servic



## dations

with principles in the Canada Health Act, publicly all Canadians, including individuals living with an nave equitable access to medically necessary all Canadians – no matter where they live and scrimination.

nd and implement specialized services to address all levels of treatment intensity and for patients of everities;

pread of leading practices aimed at better matching nt to patient needs (e.g. Short Treatment Allocation Ds (STATED))<sup>xxix</sup>.

ommunication and sharing of best practices and based care across professionals, providers / health ocial services.

tworked services for EDs across Canada based on and evidence-based practices.

networks in provinces and territories where they st to coordinate resources and collaborative care.

pan-Canadian 'network of networks' to link hese networks across provinces and territories support dissemination of best practice and on of care.

Task Force to improve transitions from youth to ces.

# 3. Treatment Recommendations (con't)

Objectives	Recommendations		
<b>3.1</b> To improve access to appropriate care in Canada;	20. Develop stronger partnerships with primary care to ensure that individuals with EDs are identified early.		
	21. Develop bilingual, culturally appropriate and gender- sensitive <sup>xxx</sup> services for marginalized sub-populations.		
in continuum of care for	22. Develop and deliver training in sensitivities regarding EDs in non-ED treatment environments (e.g. obesity clinics).		
<b>3.4</b> To support outcomes evaluation of care and the development of indicators of recovery.	23. Encourage provincial health authorities to support participation of Eating Disorder clinicians in learning opportunities to enhance their treatment competencies (e.g. EDAC-ATAC conference).		
	24. Develop and deliver specialized training to optimize treatment planning for and the integration of treatment(s) that addresses complex comorbidities.		

# 4. Caregiver Support Recommendations

Objectives	Recommend
<b>4.1</b> To connect caregivers with evidence-based and	25. Engage car recommen
evidence-informed resources and information to • help them better navigate health care system and social	26. Validate an how this St of recomm Comprehe of Adults w
<ul> <li>services;</li> <li>support their own health;</li> <li>support the recovery of the person they are caring for; and</li> <li>4.2 To develop and</li> </ul>	27. Continue to Caregiver d caregiver c caregivers' illness, invo navigating engageme
propose improvements to existing health, workplace and social policies and programs so that they are more responsive to caregiver needs.	28. Develop, w new or ada programs t need respi for their lov
	29. Further exa effectivene Health Care with Menta
	30. Develop ne better acco need to ful
	31. Work with o shared, spr living with



## dations

aregivers formally in the design and development of ndations in this Strategy.

nd prioritize with caregivers and other stakeholders Strategy can support the implementation nendations in the National Guidelines for a ensive Service System to Support Family Caregivers with Mental Health Problems and Illnesses.

to work collaboratively with the MHCC's Family Advisory Committee and national and provincial organizations to support efforts aimed at improving s' access to resources and information about mental volvement in system and program planning; g health care systems and social services; and ent, when appropriate, as part of the care team.

with national and provincial caregiver organizations, lapted provincial and organizational policies and that will provide better support to all caregivers who ite and/or who are trying to access appropriate care oved ones.

camine with caregivers and stakeholders the value and ess of existing caregiver guides such as the Mental regiver Guide: A Guide for Caregivers of Persons Living tal Illness or Experiencing Mental Health Challenges<sup>xii</sup>.

new or adapt existing workplace policies that can commodate periodic absences from work due to a ulfill caregiver responsibilities.

employers to identify better practices that can be read and adapted in support of caregivers of people mental illness.

# 5. Training Recommendations

Objectives	Recommendations
<ul> <li>5.1 To identify post graduate and continuing professional development training pathways for EDs;</li> <li>5.2 To develop clinical competencies related to the provision of health services to individuals with EDs and to their families; and</li> </ul>	32. Coordinate and disseminate training in evidence-based treatments as they become available for current professionals.
	33. Develop a continuing professional development webinar series by Canadian ED specialists with appropriately recognized learning credits.
	34. Deliver training in sensitivities regarding EDs in non-ED treatment environments, which could include correctional settings and addictions.
<b>5.3</b> To encourage future graduates to pursue research careers in the area of EDs.	35. Disseminate best practices via Telehealth for rural and remote areas.
	36. Enhance training in Family Practice settings, primary care networks/health teams and integrated health networks to support ED screening and early identification.
	37. Support professionals wishing to transition their practice into the field of EDs via intensive exposure within specialized ED settings.
	38. Identify training/exposure opportunities to other healthcare professionals who come into contact with individuals with EDs (e.g., Gastroenterologists, Emergency Department staff, Paramedics, etc.).
	39. Create a task force to identify minimum standards for graduate education in EDs within all health disciplines (including development of a standard curriculum for EDs).

# 6. Research Recommendations

Objectives	Recommend
<b>6.1</b> To identify research priorities, needs and gaps in the field of EDs in Canada;	40. Continue to with lived ex in the field s those affect these priorit
<b>6.2</b> To enhance Canadian collaborative research and support for EDs research in Canada;	41. Support a sy priorities ac
and	42. Update <sup>xlviii</sup> e
<b>6.3</b> To enhance ongoing surveillance, performance measures and data related to EDs.	43. Update exis they include
	44. Enhance up knowledge
	45. Identify/creatistic disciplines.
	46. Encourage c objectives a CIHR, SSHRC
	47. Establish a C
	48. Establish a C the EDs Cen
	49. Increase fina
	50. Create a nat capabilities



## dations

to identify research needs and priorities by people experience, caregivers, and health care professionals I so that research questions of most importance to cted by an ED are carried out and widely promote rities<sup>xiii</sup>.

systematic approach to the identification of research across the spectrum of EDs and affected populations.

existing EDs data and relevant indicators in Canada.

isting mental health performance measures so that de EDs<sup>xlix</sup>.

ptake of available ED research by using various e translation strategies and models.

eate research training opportunities across all

collaboration, organization and planning of research and combination of expertise to attain federal (i.e., RC) funding to carry out pan-Canadian research.

Canadian Research Chair for EDs.

Canadian Research Network for EDs, modelled on entres for Excellence in British Columbia.

nancial support for research related to EDs.

ational data repository for enhanced research s that reflect the Canadian context.

# **Endnotes**

- See notably Canadian Paediatric Surveillance Program, Canadian Paediatric Society (2018), CPSP 2017 Results, https://www.cpsp.cps.ca/uploads/publications/CPSP-2017-Results 1.pdf.
- ii Birmingham, C.L., Su, J., Hlynsky, J.A., Goldner, E.M. & Gao, M. (2005), "The mortality rate from anorexia nervosa," International Journal of Eating Disorders, 38(2), pp. 143-6; Girz L, Lafrance Robinson A, Tessier C. (2014), "Is the next generation of physicians adequately prepared to diagnose and treat eating disorders in children and adolescents?" Eating Disorders: Journal of Treatment & Prevention, 22(5), pp. 375-85.
- iii Hart, L. M., Granillo, M. T., Jorm, A. F., & Paxton, S. J. (2011), "Unmet need for treatment in the eating disorders: A systematic review of eating disorder specific treatment seeking among community cases," Clinical Psychology Review, 31(5), pp. 727-735. http://doi. org/10.1016/j.cpr.2011.03.004; and Mitchison, D., & Hay, P. J. (2014), "The epidemiology of eating disorders: Genetic, environmental, and societal factors," Clinical Epidemiology, p. 90, http://doi.org/10.2147/CLEP.S40841.
- iv Statistics Canada (2018), Death, causes of death and life expectancy, 2016, Table 13-10-0801-01 (formerly CANSIM 102-0564).
- Arcelus, J., Mitchell, A.J., Wales, J., & Nielsen, S. (2011), "Mortality rates in patients with v anorexia nervosa and other eating disorders: A meta-analysis of 36 studies," Archives of General Psychiatry, 68, pp. 724-731.
- Berkman, N.D., Lohr, K.N., & Bulik, C.M. (2007), "Outcomes of eating disorders: A systematic vi review of the literature," International Journal of Eating Disorders, 40, pp. 293-309.
- Arcelus, J., Mitchell, A.J., Wales, J., & Nielsen, S. (2011), "Mortality rates in patients with vii anorexia nervosa and other eating disorders: A meta-analysis of 36 studies."
- Smink, F.E., van Hoeken, D., & Hoek, H.W. (2012), "Epidemiology of eating disorders: viii Incidence, prevalence and mortality rates," Current Psychiatry Reports, 14, pp. 406-414

- ix org/10.1111/josh.12590.
- Х Public Health, 109, p. 61, https://doi.org/10.17269/s41997-018-0025-x
- xi
- xii
- xiii details.html?applId=363895&lang=en.
- xiv 0501-4.
- XV
- Ibid. xvi

See, notably, Roberts, K.C. (2012), "Overweight and obesity in children and adolescents: results from the 2009 to 2011 Canadian Health Measures Survey," Health Rep. 23(3), pp. 37-41; and Lillico, H.G. et al. (2014), "The prevalence of eating behaviors among Canadian youth using cross-sectional school-based surveys," BMC Public Health. 14, p. 323; See also, Luongo, N.M. (2018), "Disappearing in plain sight: An exploratory study of co-occurring eating and substance abuse dis/orders among homeless youth in Vancouver, Canada," Women's Studies International Forum, 67, pp. 38-44, https://doi.org/10.1016/j. wsif.2018.01.003; and Godin, K.M., Patte, K.A. and Leatherdale, S.T. (2018), "Examining Predictors of Breakfast Skipping and Breakfast Program Use Among Secondary School Students in the COMPASS Study" Journal of School Health, 88, pp. 150-158, https://doi.

Raffoul, A., Leatherdale, S.T. & Kirkpatrick, S.I. (2018), "Dieting predicts engagement in multiple risky behaviours among adolescent Canadian girls: a longitudinal analysis" Can J

McVey G., Tweed S. & Blackmore E. (2004), "Dieting among preadolescent and young adolescent females," CMAJ Canadian Medical Association Journal, 170(10), pp. 1559-61; McVey, G.L., Tweed, S., & Blackmore, E. (2005), "Correlates of dieting and muscle gaining behaviors in 10-14 year-old males and females. Preventive Medicine, 40(1), 1-9; Bernier, C.D., Kozyrskyj, A., Benoit, C., Becker, A.B. & Machessault, G. (2010), "Body image and dieting attitudes among preadolescents," Canadian Journal of Dietetic Practice and Research, 71(3), pp. e34-e40; and Gusella, J., Goodwin, J. & van Roosmalen, E. (2008), "'I want to lose weight': Early risk for disordered eating?" Paediatric Child Health, 13(2), pp. 105-110.

Pinhas, L., Morris, A. Crosby, R.D., & Katzman, D.K. (2011), "Incidence and age-specific presentation of restrictive Eating Disorders in children. A Canadian paediatric surveillance program study," Archives of Pediatrics and Adolescent Medicine, 165, pp. 895-899a.

Obeid, N. McVey, G.L., Norris, M.L. and Preskow, W.E. (2017), Building capacity for patientoriented research in eating disorders, http://webapps.cihr-irsc.gc.ca/decisions/p/project\_

See Hunt *et al.*, "The role of boundary spanners in delivering collaborative care: a process evaluation," BMC Family Practice (2016), 17, p. 96, https://doi.org/10.1186/s12875-016-

Salas et al. (2017), "A critical analysis of obesity prevention policies and strategies," Can J Public Health, 108(5-6), pp. e598-e608, https://doi.org/10.17269/CJPH.108.6044.

- Dimitropoulos, G. et al. (2015), "'You don't have anorexia, you just want to look like a xvii celebrity': perceived stigma in individuals with anorexia nervosa," Journal of Mental Health, 25(1), pp. 1-8.
- xviii https://www.ipsos.com/en-ca/half-50-canadians-have-been-exposed-eating-disorderswhether-its-someone-they-know-or-themselves
- Mond, J. M., Hay, P. J., Paxton, S. J., Rodgers, B., Darby, A., Nillson, J., Quirk, F., & Owen, C. xix (2010), "Eating disorders 'mental health literacy' in low risk, high risk and symptomatic women: Implications for health promotion programs," Eating Disorders: The Journal of Treatment & Prevention, 18(4), pp. 267-85, https://doi.org/10.1080/10640266.2010.490115.
- Canadian Teacher's Federation (2012), Understanding Teachers' Perspectives on Student XX Mental Health: Findings from a National Survey http://www.ctf-fce.ca/Research-Library/ StudentMentalHealthReport.pdf
- Carter, J. C., Stewart, D. A., Dunn, V. J., & Fairburn, C. G. (1997), "Primary prevention of eating xxi disorders: might it do more harm than good?" International Journal of EDs, 22(2), pp. 167-172.
- xxii Carter et al. (1997); Also see Yager, Z., "What not do when teaching about eating disorders," Journal of the HEIA, 13(1).
- Chiodo, J. & Latimer, P. R. (1983), "Vomiting as a learned weight-control technique in bulimia," xxiii Journal of Behavior Therapy and Experimental Psychiatry, 14(2), pp. 131-135; Fairburn, C. G. & Cooper, P. J. (1982), "Self-induced vomiting and bulimia nervosa: an undetected problem," British Medical Journal, 284(6323), pp. 1153-1155; O'Dea, J. (2000), "School-based interventions to prevent eating problems: First do no harm," EDs: The Journal of Treatment & Prevention, 8(2), pp. 123-130, https://doi.org/10.1080/10640260008251219; Thomsen, S. R., Kelly McCoy, J., & Williams, M. (2001), "Internalizing the impossible: Anorexic outpatients' experiences with women's beauty and fashion magazines," EDs: The Journal of Treatment & Prevention, 9(1), p. 57, https://doi.org/10.1080/106402601300187731.
- xxiv Pinhas, L., Wong, J., & Woodside, D. B. (2014), "Early Intervention in EDs," in P. Byrne and A. Rosen (Eds.), Early Intervention in Psychiatry: El of nearly everything for better mental health, John Wiley & Sons.
- Waller, G., D'Souza Walsh, K., & Wright, C. (2015), "Impact of education on clinicians' attitudes XXV to exposure therapy for EDs.," Behavior Research and Therapy, pp. 76-80. https://doi. org/10.1016/i.brat.2015.11.014.

- xxvi Challenges," Minerva Psichiatr., 51(3), pp. 207–216.
- xxvii inpatients with eating disorders," Psychosom Med., 68(3), pp. 454-62.
- xxviii 77; also see examples from New Brunswick and Ontario:
  - help-1.4140954;
  - business-are-scrambling-to-react.html;
  - local-news/mental-health-funding;

  - increasing-access-to-mental-health-care-in-north-simcoe.html;
- treatment-allocation-tool-for-eating-disorders-stated/



See notably, Lock, J. (2010), "Treatment of Adolescent Eating Disorders: Progress and

Blinder B.J., Cumella E.J. and Sanathara V.A., (2006), "Psychiatric comorbidities of female

See notably, Smith D.H. and Hadorn D.C. (2002), "Lining up for children's mental health services: a tool for prioritizing waiting lists," J Am Acad Child Adolesc Psychiatry, 41, pp. 367-

a. Betts, S., "Young Canadians are reaching out for mental health help more than ever, investigation shows," CBC News. June 1, 2017. Accessed September 10, 2019. http://www.cbc.ca/news/canada/new-brunswick/young-people-mental-health-

b. Cribb, R. Ovid, N., Lao, D. and Bigham, B., "Demand for youth mental health services is exploding. How universities and business are scrambling to react," The Toronto Star. May 29, 2017. Accessed September 10, 2019. https://www.thestar.com/news/ canada/2017/05/29/youth-mental-health-demand-is-exploding-how-universities-and-

c. Cobb, C. "As demand for mental health services grows, funding stagnates," The Ottawa Citizen. April 1, 2015. Accessed September 10, 2019. http://ottawacitizen.com/news/

d. Children's Mental Health Ontario (2017), Ontario Budget 2017: Wynne Budget Ignores Children's Mental Health Crisis. Accessed September 10, 2019. https://cmho.org/news/ 6519757-ontario-budget-2017-wynne-budget-ignores-children-s-mental-health-crisis;

e. Government of Ontario (2017), Ontario Increasing Access to Mental Health Care in North Simcoe. Accessed September 2010. https://news.ontario.ca/mohltc/en/2017/10/ontario-

f. Canadian Broadcasting Corporation, "CHEO, Royal wait lists for youth mental health treatment still 'extensive'" CBC News. October 29, 2015. Accessed September 10, 2019. http://www.cbc.ca/news/canada/ottawa/cheo-royal-wait-times-extensive-1.3294407

xxix Geller. J., Raudzus, J. and Yoon, P., Providence Health Care (2017), Short Treatment Allocation Tool for Eating Disorders (STATED) cited by Health Standards Organization (HSO), Leading Practices. Accessed September 10, 2019. https://healthstandards.org/leading-practice/short-

- See notably Judd, F., Armstrong, S. and Kulkarni, J. (2009), "Gender-Sensitive XXX Mental Health Care," Australasian Psychiatry, 17:2, pp. 105–111, https://doi. org/10.1080/10398560802596108 and Greenberg, S. T. and Schoen, E. G. (2008), "Males and eating disorders: Gender-based therapy for eating disorder recovery," Professional Psychology: Research and Practice, 39(4), pp. 464-471, https://doi.org/10.1037/0735-7028.39.4.464. Related to future training, see also Calzo J.P., Katz-Wise S.L., Charlton, B.M., Gordon, A.R. and Krieger, N. (2019), "Addressing the dearth of critical gender analysis in public health and medical pedagogy: an interdisciplinary seminar to generate student-created teaching examples", Critical Public Health, 29:1, pp. 18-26, https://doi.org/10.1080/09581596.2017.1419166.
- See British Columbia, Clinical Practice Guidelines for the BC Eating Disorders Continuum xxxi of Services (2012); American Psychiatric Association (2006), Practice Guideline for the Treatment of Patients with Eating Disorders, Third Edition, https://psychiatryonline. org/guidelines and National Institute for Health and Care Excellence (2017), Eating disorders: recognition and treatment, NG 69, https://www.nice.org.uk/guidance/ng69.
- Mental Health Commission of Canada (MHCC), Caregiving, https://www. xxxii mentalhealthcommission.ca/English/what-we-do/caregiving
- xxxiii Ibid.
- xxxiv Although please see Fjermestad, K. W., Rø, A. E., Espeland, K. E., Halvorsen, M.S. and Halvorsen, I. M. (2019) "'Do I exist in this world, really, or is it just her?'Youths' perspectives of living with a sibling with anorexia nervosa," Eating Disorders, https://doi. org/10.1080/10640266.2019.1573046 and Elufong, P.R. and Conti, J. (2019), "We don't really know what else we can do": Parent experiences when adolescent distress persists after the Maudsley and family-based therapies for anorexia nervosa, Journal of Eating Disorders, 7:5, https://doi.org/10.1186/s40337-019-0235-5.
- See notably, MHCC, National Guidelines for a Comprehensive Service System to Support XXXV Family Caregivers of Adults with Mental Health Problems and Illnesses, pp. 8, 11-14, https:// www.mentalhealthcommission.ca/English/document/8601/national-guidelinescomprehensive-service-system-support-family-caregivers-adults-ment
- xxxvi NIED (2018), NIED 2018 Caregiver's Need Assessment Survey.
- xxxvii Statistics Canada (2007, 2012), General Social Survey, https://www150.statcan.gc.ca/n1/ pub/89f0115x/89f0115x2013001-eng.htm.

- xl of Adults with Mental Health Problems and Illnesses
- xli *Ibid.*, p. 32.
- xlii mental-health-caregiver-guide.aspx
- xliii treatment, NG 69, https://www.nice.org.uk/guidance/ng69.
- xliv clinical-features-and-diagnosis
- xlv fichier\_these-cadwallader-201800863.pdf
- xlvi community-care-687245651.html
- xlvii decisions/p/project details.html?applId=363895&lang=en



xxxviii Chen, W-Y., and Lukens, E. (2011), "Well being, depressive symptoms, and burden among parent and sibling caregivers of persons with severe and persistent mental illness," Social Work in Mental Health, 9, pp. 397–416, https://dx.doi.org/10.1080/15332985.2011.575712.

xxxix Ennis, E. and Bunting B.P. (2013), "Family burden, family health and personal mental health," BMC Public Health, 13:255, http://www.biomedcentral.com/1471-2458/13/255.

MHCC, National Guidelines for a Comprehensive Service System to Support Family Caregivers

Ottawa Public Health (2015), http://www.ottawapublichealth.ca/en/public-health-topics/

American Psychiatric Association (2006), Practice Guideline for the Treatment of Patients with Eating Disorders, Third Edition, https://psychiatryonline.org/guidelines and National Institute for Health and Care Excellence (2017), Eating disorders: recognition and

https://www.uptodate.com/contents/eating-disorders-overview-of-epidemiology-

Cadwallader, J.S., TCA-MG : Place du médecin généraliste dans la prise en charge des patients atteints de troubles des conduites alimentaires en France, Doctoral Thesis, l'Université Paris-Saclay, 2018, p. 24, http://www.sfmg.org/data/generateur/generateur fiche/1398/

Canadian Institute for Health Information (2018), A Canadian first: CIHI to measure access to mental health and addictions services and to home and community care. Accessed September 10, 2019. https://www.newswire.ca/news-releases/a-canadian-first-cihito-measure-access-to-mental-health-and-addictions-services-and-to-home-and-

For overview, see Obeid, N., McVey, G.L., Norris, M.L. and Preskow, W.E. (2017), Building capacity for patient-oriented research in eating disorders, http://webapps.cihr-irsc.gc.ca/

xlviii Government of Canada (2019). Mental health and wellness. Accessed September 10, 2019. https://www.canada.ca/en/public-health/topics/mental-health-wellness.html

- Goffin, P., "Cross-country mental health data should be standardized, researchers say," xlix The Toronto Star, August 30, 2017. Accessed September 10, 2019. https://www.thestar. com/news/gta/2017/08/30/cross-country-mental-health-data-should-be-standardizedresearchers-say.html
- Canadian Institute for Health Information (2019). Mental health and addictions. Accessed September 10, 2019. https://www.cihi.ca/en/mental-health-and-addictions
- li Standing Committee on the Status of Women (2014), Eating Disorders among Girls and Women in Canada. Accessed September 10, 2019. https://www.ourcommons.ca/ DocumentViewer/en/41-2/FEWO/report-4.
- Hoek, H., (2016), "Review of the worldwide epidemiology of eating disorders," Current lii Opinion in Psychiatry, 29(1), https://doi.org/10.1097/YCO.00000000000282.
- liii World Health Organization (2011), "WHO highlights global underinvestment in mental health care." Accessed September 10, 2019. http://www.who.int/mediacentre/news/ notes/2011/mental health 20111007/en/
- NIED (2018), NIED 2018 Caregiver's Need Assessment Survey. liv
- Government of Canada, Federal Framework for Suicide Prevention Act, S.C. 2012, c. 30. lv Accessed September 10, 2019. http://laws.justice.gc.ca/eng/AnnualStatutes/2012\_30/.
- lvi Williams M. and Leichner, P. (2006), "More training needed in eating disorders: A time cohort comparison of Canadian psychiatry residents,". Eating Disorders: Journal of *Treatment* & *Prevention*, 14(4), pp. 323-34.
- Su, J.C. and Birmingham, C.L. (2003), "Anorexia nervosa: The cost of long-term disability," lvii *Eating and Weight Disorders*, 8(1), pp. 76-9.
- For example, see de Oliviera et al. (2016), "Cost evaluation of out-of-country care for lviii patients with eating disorders in Ontario: a population-based study," CMAJO, 4 (4), pp. e661 - e667, https://doi.org/10.9778/cmajo.20160057.
- lix Su, J.C. and Birmingham, C.L. (2003), "Anorexia nervosa: The cost of long-term disability," Eating and Weight Disorders.
- The Butterfly Foundation (2012), Paying the price: The economic and social impacts of lх eating disorders in Australia.

lxi Ibid.

- lxii e71, https://doi.org/10.1503/cmaj.170132.
- lxiii research in eating disorders.
- lxiv caregivers.html.
- lxv



Dutton, D.J. et al. (2018), "Effect of provincial spending on social services and health care on health outcomes in Canada: an observational longitudinal study," 190 (3), pp. e66 -

Obeid, N. McVey, G.L., Norris, M.L. and Preskow, W.E., Building capacity for patient-oriented

Public Health Agency of Canada (2015), Eating disorders in teens: Information for parents and caregivers. Accessed September 10, 2019. https://www.canada.ca/en/public-health/ services/publications/healthy-living/eating-disorders-teens-information-parents-

formerly the Canadian Association of Pediatric Healthcare Centres (CAPHC). Accessed September 10, 2019. https://ken.childrenshealthcarecanada.ca/xwiki/bin/view/Main/ Canadian Eating Disorders Strategy: 2019-2029